

Project Summary for IgANN website

Project Title

Follow-up microscopic hematuria as a risk of progression in adults and children with IgA vasculitis nephritis (IgAVN).

Primary Investigators

Leader: Ladan Zand, Nephrology and Hypertension, Medicine, Mayo Clinic, Rochester, Minnesota, USA
In collaboration with Fernando Fervenza, Division of Nephrology and Hypertension, Department of Medicine, Mayo Clinic, Rochester, Minnesota, USA.

Rosanna Coppo, Fondazione Ricerca Molinette, Regina Margherita Hospital, Turin, Italy

Statistical analysis:

Lisa Vaughan Department of Biostatistics, Mayo Clinic, Rochester, Minnesota, USA.

Study group investigators and data providers:

Ladan Zand, Nephrology and Hypertension, Medicine, Mayo Clinic, Rochester, Minnesota, USA.

Maria Luisa Russo, Fondazione Ricerca Molinette, Turin, Italy

Sean Barbour, University of British Columbia, Division of Nephrology, Vancouver, Canada;

Lee Er, BC Renal, British Columbia Renal, Vancouver, Canada

Mark Haas, Department of Pathology and Laboratory Medicine, Cedars-Sinai Medical Center, Los Angeles, USA

Manuel Praga, Nephrology, Instituto de Investigación Hospital 12 de Octubre, Madrid, Spain.

Brief Description

After years of being largely overlooked, microscopic hematuria has re-emerged as an important risk factor for disease progression in primary IgAN (1), which has been confirmed by several studies (2-4). A recent review updated the evidence supporting microscopic hematuria as a biomarker of disease activity and disease progression in IgAN and suggests its use during the follow-up (5). Notably, the value of microscopic hematuria in renal non-IgA related vasculitis has also been recently reported (6). However, the value of persistent microscopic hematuria on renal outcomes in IgA vasculitis nephritis (IgAVN) has not been previously investigated. The present study was designed to address this gap in knowledge. This study is based on the database collected for the IgANN research on IgAVN (7) with additional updates. Statistical analysis will be performed at the Mayo Clinic.

A letter was sent to ensure that each center approved the use of their stored data: a positive answer was received from all the centers. Additional updates were requested and obtained, resulting in a final cohort of 493 cases of IgAVN who had available MEST-C score and longitudinal follow-up data on microscopic hematuria quantification. The IgAVN database was transferred from UBC Vancouver to the Mayo Clinic in Rochester in September 2024. Data of the existing database has been supplemented by other data provided by individual centers participating in the IgANN Group. The Statistical analysis is ongoing with exchange of information between the statistical groups.

References

- 1) Sevillano et al JASN 2017; 28: 33089.3099
- 2) Coppo R and Fervenza F JASN 2017; 28:2831-2834
- 3) Bobart S et al NDT 2021; 26:36:840-847
- 4) El Karoui K, Fervenza FC, De Vriese AS. J Am Soc Nephrol. 2024;35:103-116.
- 5) Zand L, Fervenza FC, Coppo R. Clin Kidney J. 2023 Dec 4;16(Suppl 2):ii19-ii27.
- 6) Benichou N, et al French Vasculitis Study Group (FVSG) and European Vasculitis Society (EUVAS) investigators. Kidney Int. 2023; 103:1144-1155.
- 7) Barbour SJ, et al .Clin J Am Soc Nephrol. 2024;19:438-451.